

Prevalence of Diabetic Foot Ulcers among People with Diabetes Mellitus

Daris Francis^{1*}, K. Kotteswaran¹, Pramod P. V²

¹Saveetha College of Physiotherapy, Saveetha Institute of Medical and Technical sciences,
Thandalam, Chennai, India

³Lourde Institute of Allied Health Sciences, Thaliaparamba, Kannur, India

*Email: darisfrancis@gmail.com

Abstract

Introduction: With 8.6% prevalence, India tops the list in terms of the burden of diabetes. A major complication of diabetes mellitus is diabetic foot ulcer, which affects type 2 diabetes more frequently. This study's objective was to determine the prevalence and risk variables that contribute to the development of diabetic foot ulcers.

Method: A hospital-based cross-sectional survey was conducted with 108 type 2 diabetic patients who were chosen from the outpatient department of a major diabetic care facility. The socio-demographic data was gathered. The clinical biochemistry lab attached to the hospital performed the measurements of biochemical variables. The Wagner Meggitt Classification was used to stage each patient's diabetic foot.

Results: Prevalence of Diabetic Foot Ulcer was 10.2%. Grade III was the most prevalent stage of presentation, followed by Grade IV according to classification system by Wagner Meggitt. No significant correlation between the DFU group and other metabolic parameters such as hypertension, retinopathy, neuropathy, HbA1c, cholesterol, HDL, and LDL were discovered.

Conclusions: There were 10.2% of diabetic individuals presented with foot ulcers. Diabetes related foot ulcers were more common in males with an average age of 56.08± 0.69 years.

Keywords

Diabetic foot ulcer, Type 2 diabetes mellitus, Wagner-Meggitt Classification, Diabetic Peripheral Neuropathy.

Introduction

Diabetes Mellitus is a global epidemic (IDF Diabetes Atlas Group, 2015). With a regional prevalence of 8.8%, 88 million individuals in South East Asia suffer from diabetes, 77 million of whom reside in India (IDF Atlas, 2019). Diabetes prevalence among Indians ranges from 12–17% in urban areas to 2.4% in rural areas (Bengalorkar & Nagendra, 2011). A full-thickness wound that is found at a level distal to the ankle in diabetic individuals is commonly known as a diabetic foot ulcer (Apelqvist et al., 2000; Reiber & Ledoux, 2003). According to estimates, 6.3% of people worldwide have diabetic foot ulcers. Type 2 diabetics and males are more likely to get these ulcers than females (Zhang et al., 2017). In the upcoming ten years, it's anticipated that the prevalence of diabetic foot disease would increase even more (Amin & Doupis, 2016; Boulton et al., 2005). The frequency and incidence of foot issues in India are not well understood. This study's objective was to gather information on the prevalence and risk factors for developing foot ulcers particularly those connected to diabetes.

Submission: 28 May 2023; **Acceptance:** 17 August 2023



Copyright: © 2023. All the authors listed in this paper. The distribution, reproduction, and any other usage of the content of this paper is permitted, with credit given to all the author(s) and copyright owner(s) in accordance to common academic practice. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license, as stated in the website: <https://creativecommons.org/licenses/by/4.0/>

Methodology

The outpatient surgery department of Lourde Hospital in Thaliparamba, Kerala, was the site of the hospital-based cross-sectional study. Based on the criteria for inclusion and exclusion, subjects were chosen. The study included all cases of Type 2 diabetes mellitus and that were older than 18 and had lived in the study area for at least a year. Patients with Hansen's disease, gestational diabetes mellitus, stroke, bilateral below knee amputations, and foot deformities caused by other conditions were also disqualified. Data was collected during the period of March 15 to March 25, 2023. Consecutive samples of 108 patients were chosen (Ramalingam et al., 2023).

Ethical Considerations and Permissions

All subjects voluntarily provided their informed consent. Before the investigation began, the institutional ethics committee's approval was obtained.

Outcome measures

Diabetic Foot Ulcer was the outcome measure for the study. Presence of Diabetic Foot Ulcer was taken from clinical records. Each patient's diabetic foot was staged using the Wagner Meggitt Classification of Diabetic Foot Ulcers (Shah et al., 2022). The primary investigator used the questionnaire approach to obtain the demographic data.

Statistical Analysis

SPSS (Statistical Package for Social Sciences) V.15.0 (SPSS South Asia, Bangalore) for Windows was used to tabulate and analyse the gathered data. The proportions of the findings and their 95% confidence intervals were provided. The link was discovered using the chi-square test, and a p-value of 0.05 or lower was deemed significant. In order to identify the risk factors, multiple logistic regressions were performed.

Results

Participants' socio-demographic traits

One hundred eight people in all participated in the study. Women made up more than half of the respondents (68), while men made up the remaining (40). Age 56 to 65 makes up the majority of their demographic. Nearly 41.66 percent of the study's participants had diabetes for 16 to 20 years. There was systemic hypertension in 69.44%. The majority of respondents (68.51%) had a family history of diabetes and were literate (65.74%). The most frequent consequences were reported to be neuropathy (71.29%), followed by retinopathy (21.29%) (Table 1).

Table 1. Socio-demographic variables of participants with Type 2 DM (n=108)

Variables		Frequency (%)
Gender	Male	40 (37.03)
	Female	68 (62.96)
Age (years)	18-25	2 (1.85)
	26-35	5 (4.62)
	36-45	9 (8.33)
	46-55	19 (17.59)
	56-65	37(34.25)
	66-75	29 (26.85)
	76-85	7 (6.48)

	<5	2 (1.85)
	5-10	6 (5.55)
Duration of diabetes (years)	11-15	25 (23.15)
	16-20	45 (41.66)
	>20	30 (27.77)
	Underweight	13 (12.03)
BMI (kg/m ²)	Normal	37 (34.25)
	Overweight	58 (53.70)
	HbA1c (g %) (Mean±SD)	8.08±0.88
Family history	Yes	74 (68.51)
	No	34 (31.48)
Education	Literate	71 (65.74)
	Illiterate	37 (34.25)
Hypertension	Yes	75 (69.44)
	No	33 (30.55)
Neuropathy	Yes	77 (71.29)
	No	31 (28.70)
Retinopathy	Yes	23 (21.29)
	No	85 (78.70)

Table 2. Participants reported with diabetic foot ulcer

Variables		With DFU (n=11) Frequency (%)	Without DFU (n=97) Frequency (%)	p value
Gender	Male	6 (54.54)	34 (35.05)	0.581
	Female	5 (45.45)	63 (64.94)	
Age (Mean±SD)		56.08±0.69	58.09±0.70	0.383
BMI (Mean±SD)		24.09±1.65	25.01±0.79	0.960
HbA1c (g %) (Mean±SD)		8.01±0.59	8.15±0.21	0.089
Hypertension	Yes	8 (72.72)	67 (69.07)	0.704
	No	3 (27.27)	30 (30.92)	
Cholesterol (mg/dl) (Mean±SD)		216.30±15.10	214.96±26.59	0.403
Triglycerides (mg/dl) (Mean±SD)		159.09±17.06	157.63±23.07	0.710
Low Density Lipoprotein (mg/dl) (Mean±SD)		121.53±10.75	114.80±21.71	0.153
High Density Lipoprotein (mg/dl) (Mean±SD)		60.05±11.01	58.29±9.54	0.603
Neuropathy	Yes	7 (63.63)	70 (72.16)	0.098
	No	4 (36.36)	27 (27.83)	
Retinopathy	Yes	2 (18.18)	21 (21.64)	0.259
	No	9 (81.81)	76 (78.35)	
Family History of DM	Yes	6 (54.54)	68 (70.10)	0.578
	No	5 (45.45)	29 (29.89)	

P≤0.05(significant), P≤0.01(highly significant)

The frequency of foot ulcers according to participant characteristics is shown in Table 2. Males made up more than half (54.54%) of the participants with DFU. Patients with DFU had an average age of 56.08± 0.69 and a mean BMI of 24.09± 1.65. According to the results, the mean values for HbA1c (g%), cholesterol (mg/dl), triglycerides (mg/dl), low density

lipoprotein (mg/dl), and high-density lipoprotein (mg/dl) were 8.01 ± 0.59 , 216.30 ± 15.10 , 159.09 ± 17.06 , 121.53 ± 10.75 , and 60.05 ± 11.01 , respectively. Only 18.18% of them had retinopathy; instead, more than half of them had neuropathy (63.63%), hypertension (72.72%), and other conditions. When employing Z testing, no significant correlation between the DFU group and other metabolic parameters such hypertension, retinopathy, neuropathy, HbA1c, cholesterol, HDL, and LDL were discovered (Table 2).

Table 3. Diabetic foot ulcer (DFU) classification system by Wagner Meggitt

Stages of presentation	Frequency	Percentage %
Grade 0	0	0
Grade 1	0	0
Grade II	1	9.09
Grade III	5	45.45
Grade IV	3	27.27
Grade V	2	18.18
Total	11	100

The Wagner-Meggitt classification of diabetic foot was used in this investigation. Grade III (45.45%) was the most prevalent stage of presentation, followed by Grade IV (27.27%), Grade V (18.18%), and Grade II (9.09%). (Table 3).

Table 4. Prevalence of diabetic foot ulcer based on area of infection

Area of infection	Frequency	Percentage
Plantar foot ulcer	6	5.56
Dorsal foot ulcer	3	2.78
Both plantar and dorsal	2	1.85
Total	11	10.2

Eleven individuals in the study were presented with Diabetic Foot ulcer. The prevalence of foot ulcer in diabetic patient was 10.2%. Ulceration affects the plantar aspect of the metatarsal heads, the heel (5.56%), and the dorsal region of the toes (2.78%) in individuals with DFU (10.2%) (Table 4).

Discussion

Globally, diabetes mellitus exhibits epidemic-like characteristics. Even though there have been numerous research on DFU in other nations, Kerala's population has not had its pattern of complications thoroughly examined. This study's goal was to investigate the prevalence of diabetic foot ulcers and the risk factors connected to them among diabetic patients in Kannur, Kerala. In our study, the prevalence of Diabetic Foot Ulcer was 10.2%. The prevalence of foot ulcers among diabetic people worldwide ranges from 3% to 13%, according to a comprehensive review published in 2016 (Zhang et al., 2017). In India, 4.54% of individuals with type 2 diabetes mellitus were found to have diabetic foot ulcers (Das et al., 2020). The male sex has been suggested as a risk factor for the development of diabetic foot lesions in past researches (Gadepalli et al., 2006; Raja, 2007), and the present study found that the male population was more prevalent than the female population. The average age of DFU patients

was 56.08 years, which is consistent with previous research (Al-Maskari & El-Sadig,2007; Abbott et al., 2002; Al-Rubeaan et al., 2015). Only 2 (18.18%) out of the 11 individuals experienced diabetic retinopathy alterations, which is less than the previous results (Shibu et al., 2017) in Kerala. At enrolment, the majority of these patients had poor glycaemic control, with a mean HbA1C of 8.01 ± 0.59 . These results corroborate those of (Shibu et al., 2017 and Goldin et al., 2006). One of the main risk factors for diabetes patients with foot complications was systemic hypertension (72.72%), which was similar to a prior finding (Jyothylekshmy et al., 2015) in Kerala.

The study was carried out in the hospital's OP department so that we could gather a large number of patients with various foot complications. Since the study was carried out in a secondary care facility, thus the number of patients there who had diabetic foot issues may not accurately reflect the situation.

Health care providers need to focus more on DM patients who have the established risk factors for DFU. It is advised that future research on the subject incorporate behavioural aspects such as past alcohol intake history and primary data with prospective studies. If it had been a multicenter study that lasted longer, the information might have been of significantly more value.

Conclusions

Prevalence of foot ulcer in people with diabetes is 10.2% in the present study. The study emphasises the need of proper foot hygiene, the necessity of early diabetes detection, and the necessity of ongoing monitoring of diabetic complications.

Glossary of Abbreviations

DM- Diabetes Mellitus, SD- Standard Deviation, BMI- Body Mass Index, HbA1c- Glycated Haemoglobin, ANOVA- Analysis of variance, DFD- Diabetic foot syndrome or disease, DFU- Diabetic foot ulcer, WHO- World Health Organization

Acknowledgements

This research work is presented during Stride'23 International Physiotherapy conference on April 6th and 7th and the abstract is published as conference proceedings in International Journal of Physiotherapy and Occupational therapy (IJPOT).

References

- Abbott, C. A., Carrington, A. L., Ashe, H., Bath, S., Every, L. C., Griffiths, J., Hann, A. W., Hussein, A., Jackson, N., Johnson, K. E., Ryder, C. H., Torkington, R., Van Ross, E. R. E., Whalley, A. M., Widdows, P., Williamson, S., & Boulton, A. J. M. (2002). The North-West Diabetes Foot Care Study: Incidence of, and risk factors for, new diabetic foot ulceration in a community-based patient cohort. *Diabetic Medicine*, 19(5), 377–384. <https://pubmed.ncbi.nlm.nih.gov/12027925/>
- Al-Maskari, F., & El-Sadig, M. (2007). Prevalence of risk factors for diabetic foot complications. *BMC Family Practice* 8, 1-9. <https://bmcprimcare.biomedcentral.com/articles/10.1186/1471-2296-8-59>
- Al-Rubeaan, K., Al Derwish, M., Ouizi, S., Youssef, A. M., Subhani, S. N., Ibrahim, H. M.,

- & Alamri, B. N. (2015). Diabetic foot complications and their risk factors from a large retrospective cohort study. *PLoS ONE*, 10(5), 1–17. <https://pubmed.ncbi.nlm.nih.gov/25946144/>
- Amin, N., & Doupis, J. (2016). Diabetic foot disease: From the evaluation of the “foot at risk” to the novel diabetic ulcer treatment modalities. *World Journal of Diabetes*, 7(7), 153. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4824686/>
- Apelqvist, J., Bakker, K., VanHoutum, W., Nabuurs-Franssen, M., & Schaper, N. (2000). International consensus and practical guidelines on the management and the prevention of the diabetic foot. International Working Group on the Diabetic Foot. *Diabetes Metab Res Rev*, 16(1), S84-92. <https://pubmed.ncbi.nlm.nih.gov/11054895/>
- Bengalorkar, G. M., & Nagendra, K. T. (2011). Diabetic foot infections – A review. *International Journal of Biological & Medical Research*, 2(1), 453–460. [https://web-archive.southampton.ac.uk/cogprints.org/7330/1/Culture and sensitivity pattern of micro-organism isolated from diabetic foot infections in a tertiary care hospital.pdf](https://web.archive.southampton.ac.uk/cogprints.org/7330/1/Culture%20and%20sensitivity%20pattern%20of%20micro-organism%20isolated%20from%20diabetic%20foot%20infections%20in%20a%20tertiary%20care%20hospital.pdf)
- Boulton, A., Vileikyte, L., Ragnarson-Tennvall, G., & Apelqvist, J. (2005). The global burden of diabetic foot disease. *Lancet*, 366, 1719–1724. <https://pubmed.ncbi.nlm.nih.gov/16291066/>
- Das, A., Pendsey, S., Abhyankar, M., & Malabade, R. (2020). Management of Diabetic Foot in an Indian Clinical Setup: An Opinion Survey. *Cureus*, 12(6), e8636. <https://www.cureus.com/articles/28674-management-of-diabetic-foot-in-an-indian-clinical-setup-an-opinion-survey>
- Gadepalli, R., Dhawan, B., Sreenivas, V., Kapil, A., Ammini, A. C., & Chaudhry, R. (2006). A clinico-microbiological study of diabetic foot ulcers in an Indian tertiary care hospital. *Diabetes Care*, 29(8), 1727–1732. <https://pubmed.ncbi.nlm.nih.gov/16873771/>
- Goldin, A., Beckman, J. A., Schmidt, A. M., & Creager, M. A. (2006). Advanced glycation end products: Sparking the development of diabetic vascular injury. *Circulation*, 114(6), 597–605. <https://pubmed.ncbi.nlm.nih.gov/16894049/>
- IDF Atlas. (2019). *Diabetes Atlas International Diabetes Federation*. In Brussels, Belgium: International Diabetes Federation. <https://www.idf.org/e-library/epidemiology-research/diabetes-atlas/159-idf-diabetes-atlas-ninth-edition-2019.html>
- IDF Diabetes Atlas Group. (2015). Update of mortality attributable to diabetes for the IDF Diabetes Atlas: Estimates for the year 2013. *Diabetes Research and Clinical Practice*, 109, 461–465. <https://pubmed.ncbi.nlm.nih.gov/26119773/>
- Jyothylekshmy, V., Menon, A., & Abraham, S. (2015). Epidemiology of diabetic foot complications in a podiatry clinic of a tertiary hospital in South India. *Indian Journal of Health Sciences*, 8(1), 48. [https://journals.lww.com/kleu/Fulltext/2015/08010/Epidemiology of diabetic foot complications in a.9.aspx](https://journals.lww.com/kleu/Fulltext/2015/08010/Epidemiology_of_diabetic_foot_complications_in_a.9.aspx)
- Ramalingam, V., Jagatheesan, A., & Suganthirababu, P. (Eds). (2023). *Proceedings of International Physiotherapy Conference - Stride'23 in International Journal of Physiotherapy and Occupational therapy*, 1-143. <https://ijpot.com/conference.html>
- Raja, N. (2007). Microbiology of diabetic foot infections in a teaching hospital in Malaysia: a retrospective study of 194 cases. *J Microbiol Immunol Infect*, 40(1), 39–44. <https://pubmed.ncbi.nlm.nih.gov/17332905/>
- Reiber, G., & Ledoux, W. (2003). Epidemiology of diabetic foot ulcers and amputations: Evidence for prevention. In: Williams R, Herman W, Kinmonth AL, Wareham NJ, editors. *The evidence base for diabetes care*. Chichester; Hoboken (NJ): John Wiley & Sons, Ltd; pp. 641-665.
- Shah, P., Inturi, R., Anne, D., Jadhav, D., Viswambharan, V., Khadilkar, R., Dnyanmote, A., & Shahi, S. (2022). Wagner’s Classification as a Tool for Treating Diabetic Foot Ulcers: Our Observations at a Suburban Teaching Hospital. *Cureus*, 14(1), e 21501.

- <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC8861474/>
Shibu, T., Smitha, K., Gilsa, E., & Ajith, V. (2017). Biochemical Profile in Diabetic Foot Ulcer Patients – A Descriptive Study from Kerala. *IOSR Journal of Dental and Medical Sciences*, 16(1), 57–59. <https://www.researchgate.net/publication/313289981>.
Biochemical Profile in Diabetic Foot Ulcer Patients - A Descriptive Study from Kerala
Zhang, P., Lu, J., Jing, Y., Tang, S., Zhu, D., & Bi, Y. (2017). Global epidemiology of diabetic foot ulceration: a systematic review and meta-analysis†. *Annals of Medicine*, 49(2), 106–116. <https://pubmed.ncbi.nlm.nih.gov/27585063/>