

Association of Musculoskeletal Pain with Poor Quality of Sleep Among E-Gamers in a Private University in Malaysia

Leong Yi Zuang¹, Sumedha^{2*}, Jim Brown Clements¹

¹Faculty of Health & Life Sciences, INTI International University, Persiaran Perdana BBN, Putra Nilai, 71800 Nilai, Negeri Sembilan, Malaysia.

² Centre for Physiotherapy and Rehabilitation Sciences, Jamia Millia Islamia, FET - UTH Path, Ghaffar Manzil Colony, Jamia Nagar, Okhla, New Delhi 110025, India

***Email:** sumedha.svn@gmail.com, jimbrown.clements@newinti.edu.my

Abstract

Background: Electronic sports (e-sports) requires prolonged sitting and repetitive movements which makes e-gamers susceptible to develop musculoskeletal pain in different body parts. Moreover, their quality of sleep may also be affected. It is very important for physiotherapist to identify the relationship between musculoskeletal pain and poor quality of sleep among e-gamers, so as to develop more precise treatment plan and provide higher quality of patient education. **Objective:** The purpose of this study was to analyse the association between musculoskeletal pain with poor quality of sleep among e-gamers in a private university in Malaysia.

Methodology: The sample consisted of 42 e-gamers from a private university in Malaysia. A cross-sectional study was conducted by using Nordic Musculoskeletal Questionnaire and Sleep Quality Scale. The association between musculoskeletal pain and poor sleep quality was evaluated using chi-square test.

Results: The results showed that more than half of the study population reported with a higher prevalence of neck pain in the past year and in the previous week, followed by shoulders, wrists and hands. Besides, half of the participants also reported that they had very poor sleep quality (50%), followed by the category of poor quality of sleep (23.8%). E-gamers with poor sleep quality showed significant association with musculoskeletal pain in neck ($p=0.004$), shoulder ($p=0.052$), upper back ($p=0.043$), wrist and hands ($p=0.004$).

Conclusion: Our findings revealed that musculoskeletal pain in neck, shoulder, upper back, wrist and hands was significantly associated with poor sleep quality among e-gamers. Health promotion actions that contribute towards improvement in quality of sleep and prevention of musculoskeletal pain should be considered, so the performance and quality of life among the e-gamers could be improved.

Keywords

Musculoskeletal pain, Sleep quality, E-gamers, E-sports

Submission: 9 December 2021; **Acceptance:** 30 December 2021



Copyright: © 2022. All the authors listed in this paper. The distribution, reproduction, and any other usage of the content of this paper is permitted, with credit given to all the author(s) and copyright owner(s) in accordance with common academic practice. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license, as stated in the website: <https://creativecommons.org/licenses/by/4.0/>

Introduction

Electronic sports (e-sport) are a type of professional or competitive video gaming which has become a trend and a fundamental element in today's digital youth culture (Wagner, 2006). As DiFrancisco et al. (2019) have shown in their study, in US, over the past 3 years, varsity e-gaming teams have been established by more than 50 colleges. In order to improve their skills to competitive level, e-gamers are required to train for many hours per day (DiFrancisco et al., 2019). According to a study done by Kari and Karhulahti (2016), for the professional e-gamers, an average of 3.4 to 5.2 hours per day is needed for practicing, corresponding to 25 to 35 hours of egaming per week. Before a competition, the average time spent on practicing may increase up to 10 hours per day. Moreover, for most of the tournaments, up to three hours non-stop, intensive gaming is considered as normal (DiFrancisco et al., 2019). Several health-related problems, such as musculoskeletal pain, headache, sleep disorder, anxiety, stress, and decreased physical activity level have been associated with the prolonged usage of electronic devices (Dumith et al., 2010; Milde et al., 2010; Hakala et al., 2012).

DiFrancisco et al. (2019) have shown in their study, that majority of e-gamers have lack of awareness regarding good posture during practicing. In regards to the physical demands of egaming, it has been shown that the nature of this activity is similar to that of office-based work, which is prolonged sitting (Jensen et al., 2002). E-gamers tend to sit in front of a computer, laptop or phone screen, with the same position for hours at a time (Chung et al., 2019; DiFrancisco et al., 2019). Moreover, e-sport is also an activity that requires repetitive small-muscle movements specially of upper limb. According to a survey that was undertaken by Lindberg et al. (2020), 42% of Danish eSports players, taking part in structured e-sports, suffered from musculoskeletal pain. A study by DiFrancisco et al. (2019) showed that approximately 2 out of 5 eSports players, corresponding to 40% of them, suffered from either neck or back pain.

Furthermore, in a study done by Bonnar et al. (2019), game genre and gaming culture have been proved to affect the quality of sleep among e-gamers via cognitive and behavioral pathways. In order to achieve the competitive success and reaching the elite levels, e-gamers may be exposed to unhealthy wake behaviors that compromise sleep. Moreover, poor sleep behavior will be perpetuated if an e-gamer stay in a gaming team with everyone holding the similar beliefs. Light sleep and poor sleep continuity are shown to be common among e-gamers due to their low intensity physical movement (Suppiah et al., 2015). According to a study conducted by Weaver et al. (2010), an increase of 3 to 5 minutes of sleep onset latency can be observed in the subjects who had been exposed to violent videogames for 50 minutes before going to bed.

From previous studies, it has been shown that musculoskeletal pain is associated with poor quality of sleep. Kundermann et al. (2004) stated that neurophysiological changes which alter pain thresholds will be affected by sleep disturbance. Paananen et al. (2010) have shown in their study that the development of musculoskeletal pain is associated with the sleep related problems among adolescents. Moreover, compared to healthy adolescents who have no musculoskeletal pain, the adolescents who suffered with chronic musculoskeletal pain in any part of their body would have higher chances of getting poor quality of sleep (Palermo et al., 2012).

E-gamers are more susceptible to develop musculoskeletal pain in different body parts such as neck, shoulder, wrist, back and lower limbs. Their quality of sleep may also be affected. It is of utmost importance to identify the relationship between musculoskeletal pain and poor quality of sleep among e-gamers, so that preventive measures and precise treatment plan can be undertaken. Furthermore, the results of this study can act as the stepping stone to raise awareness regarding harmful effects brought about by prolonged e-gaming. Therefore, this study is conducted to identify the association of musculoskeletal pain with poor quality of sleep among e-gamers in a private university in Nilai.

Methodology

This cross-sectional study was conducted among e-gamers from a private university, Nilai, using purposive sampling method. E-gamers between age group of 18-25 years old, who spent more than 7 hours per week playing video games and have any musculoskeletal pain were recruited for this study. However, those who were unable to read English; had any sleep disorders, any mental disorders that may affect quality of sleep, any medical condition that may affect quality of sleep, or had tremendous stress were excluded from this study. The sample size was calculated using the G*Power Version 3.1.9.7. Sample size was estimated to be 37 by taking α error as 0.05, power as 0.90 and effect size as 0.5 (Cohen, 1988; 1992). The participation in the study was voluntary and withdrawal at any time was allowed. Informed consent was obtained from each participant before data collection. Ethical approval to conduct this study was obtained from the Research and Ethics

Committee of INTI International University (references no: INTI-IU/FHLSRC/BPHTI/7NY12020/013).

Nordic Musculoskeletal Questionnaire (NMQ)

Reliability study of NMQ using test-retest methodology showed that the number of different answers ranged from 0 to 23%. A range of 0 to 20% disagreement was found during the test of validity (test against clinical history and NMQ). It was acceptable in a screening tool as concluded by the authors. (Crawford, 2007).

Sleep Quality Scale (SQS)

This questionnaire has an internal consistency of 0.92 and a test-retest reliability of 0.81. The SQS is also strongly correlated with the results collected from the Pittsburgh Sleep Quality Index (Shahid et al., 2011). Using a four-point Likert-type scale, to elicit how frequently the respondents are having certain sleep behaviors (0 = "few," 1 = "some-times," 2 = "often," and 3 = "almost always"). Scores on items belong to factors 2 and 5 (restoration after sleep and satisfaction with sleep) and are reversed before being tallied. Total scores range from 0 to 84, with the higher scores indicating more acute sleep problems. (Shahid et al., 2011)

Statistical Analysis

The data was analyzed using the Statistical Package for the Social Sciences (SPSS) software Version 26. Chi-squared tests was used to identify the association between musculoskeletal pain and poor quality of sleep. The level of significance was taken as $p < 0.05$.

Results

The prevalence of neck pain was the highest in the past 12 months and in the previous week (85.7% and 66.7%, respectively), followed by shoulders pain (78.6% and 59.5%, respectively), wrists/hands pain (61.9% and 50.0%, respectively), upper back pain (45.2% and 26.2%, respectively), lower back pain (42.9% and 23.8%, respectively), knees pain (16.7% and 4.8%, respectively), elbows pain (14.3% and 9.5%, respectively), hips/thighs pain (14.3% and 4.8%, respectively) and ankles/feet pain (9.5% and 4.8%, respectively).

Based on the categorization of SQS scores shown in Table 1 (Ningrum and Kusumaningrum, 2021), out of 42 subjects, 4 (9.5%) subjects were included in the “very good” category, 2 (4.8%) subjects were included in the “good” category, 5 (11.9%) subjects were included in the “average” category, 10 (23.8%) subjects were included in the “poor” category and 21 (50%) subjects were included in the “very poor” category.

Table 1. Categorization for SQS scores

Sleep Quality	Normalization Score	Frequency	%
Very good	$X < 22$	4	9.5
Good	$22 \leq X < 27$	2	4.8
Average	$27 \leq X < 31$	5	11.9
Poor	$31 \leq X < 37.8$	10	23.8
Very poor	$X \geq 37.8$	21	50

SQS - Sleep quality scale

As shown in Table 2, there was significant association found between musculoskeletal pain in neck ($p=0.004$); shoulders ($p=0.052$); upper back ($p=0.043$); wrists/hands ($p=0.004$) and poor quality of sleep. There was no significant association found between musculoskeletal pain in elbow ($p=0.657$), lower back ($p=0.125$), hips/thighs ($p=0.885$), knees ($p=0.885$), ankles ($p=0.885$) and poor quality of sleep among the e-gamers.

Table 2. Association between musculoskeletal pain and poor quality of sleep

Body regions	Sample size (N)	Degree of value freedom (df)	Degree of value	Chi-square test p -value
Neck	42	4	15.664	0.004
Shoulders	42	4	9.408	0.052
Upper back	42	4	9.878	0.043
Elbows	42	4	2.432	0.657
Wrists/hands	42	4	15.257	0.004

Lower back	42	4	7.219	0.125
Hips/thighs	42	4	1.155	0.885
Knees	42	4	1.155	0.885
Ankles/feet	42	4	1.155	0.885

Discussions

The present study examined the association between musculoskeletal pain and poor quality of sleep among e-gamers through a self-reported survey. This study showed highest prevalence of musculoskeletal pain in neck in the past year and in the previous week, followed by shoulders, wrists and hands. Out of the 42 participants, 45.2% of the e-gamers reported that they had upper back pain in the past year, but only 26.2% of them had musculoskeletal pain in this region during the previous week. While 42.9% of the participants experienced low back pain in the past year, only 23.8% of them reported pain in this region during the previous week. The prevalence of musculoskeletal pain in this study was similar to a study undertaken by DiFrancisco et al. (2019), which showed that the highest number of complaints among e-gamers was of neck pain (41%). However, DiFrancisco et al. (2019) also reported a higher prevalence of back pain (41%) in the previous week, compared with the current study (upper back pain, 26.2%; low back pain, 23.8%). The difference could be explained by the large difference in training time between the two studies. For current study, the participants who were recruited were spending a minimum of 7 hours per week of e-gaming. However, the study carried out by DiFrancisco et al. (2019) recruited professional e-gamers who spent an average of 25-35 hours of e-gaming per week, with 50% of the participants spending more than 3 hours per day of sitting and playing without taking a break.

The results of current study showed that out of 42 participants, 50% reported to have very poor sleep quality, followed by 23.8% who reported a poor quality of sleep, while 11.9% of the egamers reported to have an average sleeping quality. A study using the same questionnaire, conducted by Ningrum and Kusumaningrum (2021) reported 24.3% of participants suffering from poor sleep quality, which was similar to the results of current study. A study with objective measures using the self-reported sleep duration by Twenge et al. (2017) showed a positive correlation between the usage of electronic devices, in term of hours, and the failure of getting 7 or more hours of sleep at night. Another study by Exelmans and Van (2015) reported that the higher the volume of video games playing, the higher the level of fatigue and insomnia. For the respondents who had e-gaming for more than 1 hour per day, their risk of having poor quality of sleep was 31% higher than normal people (Exelmans and Van, 2015). All these studies show that excessive e-gaming could affect the quality of sleep, which was similar to the results of current study.

The results of current study shows that there was association between musculoskeletal pain (neck, shoulders, upper back, wrists/hands) and poor quality of sleep among the e-gamers. This finding was in line with the previous studies done by de Souza et al. (2020), which showed that poor quality of sleep was associated with musculoskeletal pain among the public-school teachers in Brazil. Another study done by Harrison et al. (2014) also reported poor quality of sleep being associated with the presence of regional musculoskeletal pain among the adolescents in U.K. It could be explained by the finding that poor sleep quality could trigger a series of physical and emotional alterations which could further increase muscle tension and lead to musculoskeletal pain in different parts of the body (Linder et al., 2014).

Future studies should be done that includes a larger sample size, improving the external validity so that the results can be generalizable to the entire Malaysia populations. The current study has potential limitations. The questionnaires used in this study are self-reported, which may lead to the possibility of recall bias among the respondents. Besides, the target population of this study is limited to the e-gamers from private university in Nilai and cannot be generalized to entire Malaysian e-gamers fraternity.

Conclusion

This study found an association between musculoskeletal pain (neck, shoulders, upper back, wrists/hands) and poor quality of sleep among the e-gamers in private university, Nilai. Intervention studies should be carried out to test whether the treatment protocols targeting prevention or reduction of musculoskeletal pain are also effective in improving quality of sleep, and vice versa. In addition, health promotion actions that contribute towards improvement in quality of sleep and prevention of musculoskeletal pain should be considered, so the performance and quality of life among the e-gamers could be enhanced.

Acknowledgements

The authors would like to thank all the participants for their voluntary participation in this study.

References

- Bonnar, D., Lee, S., Gradisar, M., & Suh, S. (2019). Risk Factors and Sleep Intervention Considerations in Esports: A Review and Practical Guide. *Sleep Medicine Research*, 10(2), 59-66. doi: 10.17241/smr.2019.00479
- Chung, T., Sum, S., Chan, M., Lai, E., & Cheng, N. (2019). Will esports result in a higher prevalence of problematic gaming? A review of the global situation. *Journal of behavioral addictions*, 8(3), 384–394. <https://doi.org/10.1556/2006.8.2019.46>
- Cohen, J. (1988). *Statistical power analysis for the behavioral sciences* 2nd ed Hillsdale NJ Erlbaum.
- Cohen, J. (1992). A power primer. *Psychological bulletin*, 112(1), 155.
- Crawford, J. (2007). The Nordic Musculoskeletal Questionnaire. *Occupational Medicine*, 57(4), 300-301. doi: 10.1093/occmed/kqm036
- de Souza, J., Pinto, R., Tebar, W., Gil, F., Delfino, L., & Morelhão, P. et al. (2020). Association of musculoskeletal pain with poor sleep quality in public school teachers. *Work*, 65(3), 599-606. doi: 10.3233/wor-203114
- DiFrancisco-Donoghue, J., Balentine, J., Schmidt, G., & Zwibel, H. (2019). Managing the health of the eSport athlete: an integrated health management model. *BMJ Open Sport & Exercise Medicine*, 5(1), e000467. doi: 10.1136/bmjsem-2018-000467
- Dumith SC, Domingues MR, Gigante DP, Hallal PC, MenezesAM, Kohl HW. (2010). Prevalence and correlates of physical activity among adolescents from Southern Brazil. *Rev Saude*, 44:457-67
- Exelmans, L., & Van den Bulck, J. (2015). Sleep quality is negatively related to video gaming volume in adults. *Journal of sleep research*, 24(2), 189-196.
- Hakala, P. T., Saarni, L. A., Punamäki, R. L., Wallenius, M. A., Nygård, C. H., & Rimpelä, A. H. (2012). Musculoskeletal symptoms and computer use among Finnish adolescents--pain

- intensity and inconvenience to everyday life: a cross-sectional study. *BMC musculoskeletal disorders*, 13, 41. <https://doi.org/10.1186/1471-2474-13-41>
- Harrison, L., Wilson, S., & Munafò, M. (2014). Exploring the Associations between Sleep Problems and Chronic Musculoskeletal Pain in Adolescents: A Prospective Cohort Study. *Pain Research and Management*, 19(5), e139-e145. doi: 10.1155/2014/615203
- Jensen, C., Finsen, L., Sjøgaard, K., & Christensen, H. (2002). Musculoskeletal symptoms and duration of computer and mouse use. *International journal of industrial ergonomics*, 30(45), 265-275.
- Kari, T., & Karhulahti, V. M. (2016). Do e-athletes move?: a study on training and physical exercise in elite e-sports. *International Journal of Gaming and Computer-Mediated Simulations (IJGCMS)*, 8(4), 53-66.
- Kundermann, B., Krieg, J. C., Schreiber, W., & Lautenbacher, S. (2004). The effects of sleep deprivation on pain. *Pain Research and Management*, 9(1), 25-32.
- Linder, J., Jansen, G., Ekholm, K., & Ekholm, J. (2014). Relationship between sleep disturbance, pain, depression and functioning in long-term sick-listed patients experiencing difficulty in resuming work. *Journal Of Rehabilitation Medicine*, 46(8), 798-805. <https://doi.org/10.2340/16501977-1833>
- Milde-Busch, A., von Kries, R., Thomas, S., Heinrich, S., Straube, A., & Radon, K. (2010). The association between use of electronic media and prevalence of headache in adolescents: results from a population-based cross-sectional study. *BMC neurology*, 10(1), 1-10.
- Ningrum, L., & Kusumaningrum, F. (2021). Relationship Between Sleep Quality and Internet Addiction Among College Students in Yogyakarta. *Proceedings Of The 2Nd Southeast Asian Academic Forum on Sustainable Development (SEA-AFSID 2018)*. doi: 10.2991/aebmr.k.210305.
- Paananen, M. V., Auvinen, J. P., Taimela, S. P., Tammelin, T. H., Kantomaa, M. T., Ebeling, H. E., ... & Karppinen, J. I. (2010). Psychosocial, mechanical, and metabolic factors in adolescents' musculoskeletal pain in multiple locations: a cross-sectional study. *European Journal of Pain*, 14(4), 395-401.
- Palermo, T., Law, E., Churchill, S., & Walker, A. (2012). Longitudinal Course and Impact of Insomnia Symptoms in Adolescents with and without Chronic Pain. *The Journal of Pain*, 13(11), 1099-1106. doi: 10.1016/j.jpain.2012.08.003
- Shahid, A., Wilkinson, K., Marcu, S., & Shapiro, C. (2011). Sleep Quality Scale (SQS). *STOP, THAT And One Hundred Other Sleep Scales*, 345-350. doi: 10.1007/978-1-4419-98934_85
- Suppiah, H. T., Low, C. Y., & Chia, M. (2015). Effects of sports training on sleep characteristics of Asian adolescent athletes. *Biological rhythm research*, 46(4), 523-536.
- Twenge, J. M., Krizan, Z., & Hisler, G. (2017). Decreases in self-reported sleep duration among U.S. adolescents 2009-2015 and association with new media screen time. *Sleep medicine*, 39, 47-53. <https://doi.org/10.1016/j.sleep.2017.08.013>
- Wagner, M. G. (2006). On the Scientific Relevance of eSports. In *International conference on internet computing* (pp. 437-442).
- Weaver, E., Gradisar, M., Dohnt, H., Lovato, N., & Douglas, P. (2010). The effect of presleep video-game playing on adolescent sleep. *Journal of Clinical Sleep Medicine*, 6(2), 184-189.